


Electromagnetic emission from strongly interacting hadronic and partonic matter created in heavy-ion collisions



A. W. Romero Jorge^{1,2,3}, T. Song⁴, Q. Zhou⁵, E. Bratkovskaya^{2,3,4}


¹Frankfurt Institute for Advanced Studies (FIAS), ²Institute for Theoretical Physics, Goethe University, ³Helmholtz Research Academy Hessen for FAIR (HFHF), ⁴GSI Helmholtzzentrum für Schwerionenforschung, ⁵Central China Normal University, China.

1. Motivation


Dileptons serve as key probes that provide crucial insights into both the partonic and hadronic stages of heavy-ion collisions, enabling the study of in-medium effects—such as the collisional broadening of vector meson spectral functions—and the investigation of various QGP phases, thereby deepening our understanding of strongly interacting matter.

2. PHSD Transport Approach

Parton-Hadron-String Dynamics (PHSD) is a non-equilibrium microscopic transport approach that describes the full evolution of heavy-ion collisions—from initial nucleon-nucleon interactions, through QGP formation (described in terms of the DQPM model) and interactions, to hadronization and final-state interactions—using generalized off-shell transport equations based on the Kadanoff-Baym theory [1].



5. (T, μ_B) -dependent EoS on QGP dileptons



- Incorporating μ_B dependence into the QGP EoS leads to a weak impact on the total yield [3].

3. DQPM (T, μ_B)

Dynamical QuasiParticle Model (DQPM) is an effective model for the description of non-perturbative (strongly interacting) QCD based on the IQCD EoS [2].



- Quasiparticles are described in terms of dressed quark (gluon) propagators (2PI) with complex self-energies.

$$\begin{aligned} \text{gluon propagator: } & \Lambda^{-1} = P^2 - \Pi \\ \text{quark propagator: } & S_q^{-1} = P^2 - \Sigma_q \\ \text{gluon self-energy: } & \Pi = M_g^2 - i2\gamma_g\omega \\ \text{quark self-energy: } & \Sigma_q = M_q^2 - i2\gamma_q\omega \end{aligned}$$


$$\begin{cases} \text{Re } \Sigma_q : \text{ thermal masses } (M_g, M_q) \\ \text{Im } \Sigma_q : \text{ interaction widths } (\gamma_g, \gamma_q) \end{cases}$$

- Spectral functions with Lorentzian form,


$$\rho_j(\omega, \mathbf{p}) = \frac{4\omega\gamma_j}{(\omega^2 - \mathbf{p}^2 - M_j^2)^2 + 4\gamma_j^2\omega^2}$$



4. Time evolution of the fireball: ε_{tot} and μ_B




6. The fraction of energy in the QGP phase



- QGP dominates at high energy Au+Au collisions.
- QGP fraction decreases with decreasing collision energy.

7. Invariant Mass Spectra of Dielectrons in PHSD: From SIS to LHC Energies



Dileptons (e^+e^- and $\mu^+\mu^-$ pairs) are produced during the whole time evolution in heavy-ion collisions via:

Hadronic sources:

- Dalitz decays of $\pi^0, \eta, \Delta, a_1, \omega, \eta'$.
- Vector-meson decays $\rho, \omega, \phi \rightarrow e^+e^-$ (with in-medium broadening spectral functions).
- $NN \rightarrow NN e^+e^-$ and $\pi N \rightarrow \pi N e^+e^-$ bremsstrahlung (low energies).
- Correlated charm/bottom meson decays ($D\bar{D}, B\bar{B}$).

Partonic sources (QGP):

- $q\bar{q} \rightarrow e^+e^-, q\bar{q} \rightarrow g e^+e^-, qg \rightarrow q e^+e^-, \bar{q}g \rightarrow \bar{q} e^+e^-$.
- Off-shell cross sections from DQPM, including (T, μ_B) dependence.


Primary Drell-Yan (DY):

- $q\bar{q} \rightarrow e^+e^-$ in initial hard scatterings.

10. Conclusions


- Low energies (SIS):** Bremsstrahlung, Dalitz, and in-medium vector meson decays dominate, while QGP "hot spots" appear even at $\sqrt{s_{NN}} \sim 3.5$ GeV, potentially observable once Drell-Yan contribution is subtracted.
- High energies (RHIC, LHC):** Partonic processes (QGP) and heavy flavors govern the intermediate-mass region.
- Small influence of μ_B dependence of QGP EoS on dilepton mass spectra is found.
- QGP radiation overshines charm contributions at $\sqrt{s_{NN}} \simeq 25 - 30$ GeV in central Au+Au collisions, providing access to thermal QGP dileptons at BES RHIC and FAIR.

8. Excitation Function of Dielectron Yields



- Dileptons from QGP overshadow dileptons from correlated charm for $\sqrt{s_{NN}} \simeq 30$ GeV (in line with Ref. [4]).
- Subtracting the Drell-Yan contribution would allow to access the thermal QGP dileptons at $\sqrt{s_{NN}} \simeq 20$ GeV.

9. Dilepton Excess Yields



- The low-mass dilepton excess yield increases with collision energy. STAR data from [5, 6].
- The dilepton excess yield includes the ρ -meson contribution, QGP, Dalitz decays of Δ -resonances and a_1 mesons.

KEY REFERENCES

- [1] Cassing et al. (2008), PRC 78, 034919. [3] Romero Jorge et al. (2025), arXiv:2503.05253. [5] Abdulhamid et al. (2023), PRC 107, 6, L061901.
[2] Moreau et al. (2019), PRC 100, 1, 012911. [4] Song et al. (2018), PRC 97, 6, 064907. [6] STAR Coll. (2024), arXiv: 2402.01998.

ACKNOWLEDGEMENT AND PARTNERS

